



your entrance to the
knowledge resources of the
National Library of Medicine

"duroquinone" AND "ischaemia"

Search

Clear
Search

Find Terms

Limits

Results

History

Locker

Preferences

New Search

Expanded Item

Overview

What's New

Help

FAQ

Journal Citations :

Item 1 displayed (out of 3 found).

For this item only:

Download
or Display

Order
Documents

Put In
Locker

Page 1 of 3

Jump to
Page

2

Jump to
Collection

Pick A Collection

Other NLM
Resources

Ordering Info.

Clinical Alerts

ClinicalTrials.gov

HSTAT

LOCATORplus

MEDLINEplus

PubMed

TOXNET

Related
Articles

Publisher
Link

Attenuation and augmentation of ischaemia-related neuronal death by tumour necrosis factor-alpha in vitro.

Wilde GJ, Pringle AK, Sundstrom LE, Mann DA, Iannotti F.

Eur J Neurosci. 2000 Nov;12(11):3863-70.

Department of Clinical Neurological Sciences, LF73B, Level F, South Academic Block, Mailpoint 806, University of Southampton, Southampton General Hospital, Tremona Road, Southampton SO16 6YD, UK.

Upregulation of the pro-inflammatory cytokine tumour necrosis factor-alpha (TNF) occurs rapidly in the brain following ischaemia, although is unclear whether this represents a neurotoxic or neuroprotective response. We have investigated whether TNF has different actions in the pre- and postischaemic periods in a tissue culture model of cerebral ischaemia. Organotypic hippocampal slice cultures were prepared from 8-10-day-old rats and maintained in vitro for 14 days. Neuronal damage was induced by either 1 h oxygen-glucose deprivation or 3 h exposure to NMDA or the superoxide generator duroquinone, and assessed after 24 h by propidium iodide fluorescence. TNF pretreatment was neuroprotective against both oxygen-glucose deprivation and duroquinone. This effect was associated with an activation of the transcription factor NFkappaB and upregulation of manganese superoxide dismutase, and was prevented by a free radical scavenger. When addition of TNF was delayed until the postinsult period, an exacerbation of neurotoxicity occurred, which was also prevented by a free radical scavenger. The actions of TNF are determined by whether TNF is present before or after an ischaemia-related insult. Both actions are mediated through the production of free radicals, and the response to TNF is determined by whether a cell is metabolically competent to respond by synthesis of antioxidant defences.

MeSH Terms:

- o Animal
- o Benzoquinones/pharmacology
- o Cell Death
- o Free Radical Scavengers/pharmacology



Gateway

your entrance to the
knowledge resources of the
National Library of Medicine

"duroquinone" AND "ischaemia"

Search

Clear
Search

Find Terms

Limits

Results

History

Locker

Preferences

New Search

Expanded Item

Overview

What's New

Help

FAQ

Journal Citations :

Item 3 displayed (out of 3 found).

For this item only:

Download
or Display

Order
Documents

Put in
Locker

Page 3 of 3

Jump to
Page

1

Jump to
Collection

Pick A Collection

Other NLM
Resources

Ordering Info.

Clinical Alerts

ClinicalTrials.gov

HSTAT

LOCATORplus

MEDLINEplus

PubMed

TOXNET

Related
Articles

Publisher
Link

Differential vulnerability of the CA1 and CA3 subfields of the hippocampus to superoxide and hydroxyl radicals in vitro.

Wilde GJ, Pringle AK, Wright P, Iannotti F.

J Neurochem. 1997 Aug;69(2):883-6.

Department of Clinical Neurological Sciences, University of Southampton, Southampton General Hospital, England, U.K.

The relative roles of the superoxide and hydroxyl radicals in oxidative stress-induced neuronal damage were investigated using organotypic hippocampal slice cultures. Cultures exposed to 100 microM duroquinone, a superoxide-generating compound, for 3 h developed CA1-selective lesions over a period of 24 h. The damage accounted for approximately 64% of the CA1 subfield, whereas CA3 showed just 6% damage, a pattern of damage comparable to that observed following hypoxia/ischaemia. Duroquinone-induced damage was attenuated by a spin-trap agent. In contrast, hydroxyl radical-mediated damage, generated by exposure to 30 microM ferrous sulphate for 1 h, resulted in a CA3-dominant lesion. The damage developed over 24 h, similar to that observed with duroquinone, but with approximately 45% damage in CA3 compared with only 7% in CA1. These data demonstrate a selective vulnerability of the CA1 pyramidal neurones to superoxide-induced damage and suggest that of the free radicals generated following hypoxia/ischaemia, superoxide, rather than hydroxyl radical, is instrumental in producing neuronal damage.

MeSH Terms:

- o Animal
- o Benzenesulfonates/pharmacology
- o Benzoquinones/pharmacology
- o Brain Ischemia
- o Cell Death
- o Ferrous Compounds/pharmacology
- o Hippocampus/*drug effects
- o Hydroxyl Radical/ pharmacology
- o Hypoxia, Brain
- o Neurons/drug effects



Gateway

your entrance to the
knowledge resources of the
National Library of Medicine

"botulinum" AND "NG108"

Search

Clear
Search

Find Terms

Limits

Results

History

Locker

Preferences

New Search

Expanded Item

Overview

What's New

Help

FAQ

Journal Citations :

Item 3 displayed (out of 5 found).

For this item only:

Download
or Display

Order
Documents

Put in
Locker

Page 3 of 5

Jump to
Page

4

Jump to
Collection

Pick A Collection

Other NLM
Resources

Ordering Info.

Clinical Alerts

ClinicalTrials.gov

HSTAT

LOCATORplus

MEDLINEplus

PubMed

TOXNET

Related
Articles

Binding of botulinum type C1, D and E neurotoxins to neuronal cell lines and synaptosomes.

Yokosawa N, Tsuzuki K, Syuto B, Fujii N, Kimura K, Oguma K.

Toxicon. 1991;29(2):261-4.

Department of Microbiology, Sapporo Medical College, Japan.

Clostridium botulinum 125I-labeled C1 neurotoxin bound to NG108 hybridoma cell line. Unlabeled type C1 neurotoxin inhibited the binding of the labeled C1 toxin but neither types D nor E toxin. 125I-labeled type D neurotoxin bound to rat brain synaptosomes but did not bind to NG108 cells. It is suggested that receptors for types C and D or E toxin on neuronal cell membranes are different.

MeSH Terms:

- o Animal
- o Binding, Competitive
- o Botulinum Toxins/*metabolism
- o Cell Line
- o Human
- o Mice
- o Neuroblastoma/metabolism
- o Neurons/*metabolism
- o Rats
- o Synaptosomes/*metabolism
- o Tumor Cells, Cultured

Substances:

- o 0 (Botulinum Toxins)

PMID: 2048142 [PubMed - indexed for MEDLINE]

From PubMed

Page 3 of 5

Jump to
Page

4

Jump to
Collection

Pick A Collection



your entrance to the
knowledge resources of the
National Library of Medicine

"botulinum" AND "NG108"

Search

Clear
Search

Find Terms

Links

Results

History

Locker

Preferences

New Search

Expanded Item

Overview

What's New

Help

FAQ

Journal Citations :

Item 5 displayed (out of 5 found).

For this item only:

Download
or Display

Order
Documents

Put in
Locker

Page 5 of 5

Jump to
Page

1

Jump to
Collection

Pick A Collection

Other NLM
Resources

Ordering Info.

Clinical Alerts

ClinicalTrials.gov

HSTAT

LOCATORplus

MEDLINEplus

PubMed

TOXNET

Related
Articles

Binding of Clostridium botulinum type C neurotoxin to different neuroblastoma cell lines.

Yokosawa N, Kurokawa Y, Tsuzuki K, Syuto B, Fujii N, Kimura K, Oguma K.

Infect Immun. 1989 Jan;57(1):272-7.

Erratum in:

o *Infect Immun* 1989 Jul;57(7):2265. PMID: 0

Department of Microbiology, Sapporo Medical College, Japan.

Binding of type C neurotoxin (C1 toxin) from Clostridium botulinum (strain Stockholm) to neuroblastoma cell lines was studied by using biotinylated anti-toxin antibody and avidin-biotinylated peroxidase complex. The neurotoxin bound with high efficiency to mouse neuroblastoma (NS-20Y and NIE-115) cells and to hybridomas of rat glioblastoma and mouse neuroblastoma (NG108-C15) cells. The toxin bound little to human neuroblastoma, rat astrocytoma, and nonneural cell lines. Binding of the neurotoxin to NG108-C15 cells was inhibited by gangliosides (GT1b and GM1) and by monoclonal antibodies (CA-12 and C-9), although inhibition was not complete. Sequential preincubation of C1 toxin with GT1b and CA-12 caused complete inhibition. A Scatchard plot of binding of ¹²⁵I-labeled C1 toxin to NG108-C15 cells showed a hyperbolic curve. Monoclonal antibody CA-12 but not C-9 neutralized the lethal activity of the toxin toward mice. Only C-9 clearly inhibited toxin binding to GT1b. These results suggest that NG108-C15 cells have at least two kinds of receptors for C1 toxin. From the results of binding tests with neuraminidase-, pronase-, and trypsin-treated NG108-C15 cells, the chemical nature of the high-affinity site was presumed to be a glycoprotein containing sialic acid. GT1b may have an important role in low-affinity sites.

MeSH Terms:

- o Animal
- o Antibodies, Monoclonal
- o Binding, Competitive